

Quantization of Friedmann-Robertson-Walker spacetimes in the presence of a negative cosmological constant and radiation

G. A. Monerat*, E. V. Corrêa Silva†, G. Oliveira-Neto‡, ¹ L. G. Ferreira Filho§, ² and N. A. Lemos¶³

¹*Departamento de Matemática e Computação, Faculdade de Tecnologia,
Universidade do Estado do Rio de Janeiro,
Estrada Resende-Riachuelo, s/nº, Morada da Colina
CEP 27523-000, Resende-RJ, Brazil.*

²*Departamento de Mecânica e Energia, Faculdade de Tecnologia,
Universidade do Estado do Rio de Janeiro,
Estrada Resende-Riachuelo, s/nº, Morada da Colina
CEP 27523-000, Resende-RJ, Brazil.*

³*Instituto de Física, Universidade Federal Fluminense,
R. Gal. Milton Tavares de Souza s/nº, Boa Viagem
CEP 24210-340, Niterói-RJ, Brazil
(Dated: February 6, 2008)*

In the present work, we quantize three Friedmann-Robertson-Walker models in the presence of a negative cosmological constant and radiation. The models differ from each other by the constant curvature of the spatial sections, which may be positive, negative or zero. They give rise to Wheeler-DeWitt equations for the scale factor which have the form of the Schrödinger equation for the quartic anharmonic oscillator. We find their eigenvalues and eigenfunctions by using a method first developed by Chhajlany and Malnev. After that, we use the eigenfunctions in order to construct wave packets for each case and evaluate the time-dependent expected value of the scale factors. We find for all of them that the expected values of the scale factors oscillate between maximum and minimum values. Since the expectation values of the scale factors never vanish, we conclude that these models do not have singularities.

PACS numbers: 04.40.Nr, 04.60.Ds, 98.80.Qc

Keywords: quantum cosmology, Wheeler-DeWitt equation, negative cosmological constant, quartic anharmonic oscillator

* E-mail: monerat@uerj.br
† E-mail: evasquez@uerj.br
‡ E-mail: gilneto@fat.uerj.br
§ E-mail: gonzaga@fat.uerj.br
¶ E-mail: nivaldo@if.uff.br

I. INTRODUCTION

One of the motivations for the quantization of cosmological models was that of avoiding the initial *Big Bang* singularity. Since the pioneering work in quantum cosmology due to DeWitt [1], workers in this field have been attempting to prove that quantum cosmological models entail only regular space-times. An important contribution to this issue was given by Hartle and Hawking [2], who proposed the *no-boundary* boundary condition, which selects only regular space-times to contribute to the wave-function of the Universe, derived in the path integral formalism. Therefore, by construction, the *no-boundary* wave-functions are everywhere regular and predict a non-singular initial state for the Universe. Using that boundary condition, in certain particular cases the *no-boundary* wave-function can be explicitly computed [2, 3, 4]. Another way by which one may compute the wave-function of the Universe is by directly solving the Wheeler-DeWitt equation [1]. The wave-function of the Universe for some important models have been computed using this approach [5, 6]. We should mention another boundary condition, the *tunnelling* boundary condition, proposed by Vilenkin [7], to be imposed on solutions to the Wheeler-DeWitt equations. The *tunnelling* wave-function was also shown to give rise to models free from the initial *Big Bang* singularity [7]. More recently, the absence of singularities in quantum cosmological models has been investigated by using the DeBroglie-Bohm interpretation of quantum mechanics [8]. In this interpretation, one may compute the dynamical trajectories for the quantum variables of the system. In particular, since for most of the quantum cosmological models one uses minisuperspaces [9], one computes, through the DeBroglie-Bohm interpretation, the dynamical trajectories for the scale factor. Then, for the great majority of cases studied so far, the scale factor never vanishes, which implies that the model has no initial singularities [10, 11]. Although we shall restrict our attention to a model in quantum general relativity, it is important to mention that several works in loop quantum cosmology have also shown that the wave-function of the Universe is free from initial singularities [12, 13].

Several important theoretical results and predictions in quantum cosmology have been obtained with a negative cosmological constant. Considering a subset of all four-dimensional spacetimes with constant negative curvature and compact space-like hypersurfaces, Carlip and coworkers showed how to compute the sum over topologies leading to the *no-boundary* wave-function [14, 15]. These spacetimes are curved only due to the presence of a negative cosmological constant. In Ref. [14] it was shown how to obtain a vanishing cosmological constant as a prediction from the *no-boundary* wave-function and in Ref. [15] it was shown how to obtain predictions about the topology of the

Universe from the *no-boundary* wave-function. We may also mention the result in Ref. [3], where the WKB *no-boundary* wave-function of a homogeneous and isotropic Universe with a negative cosmological constant was computed. Due to the regularity condition imposed upon the space-times contributing to the *no-boundary* wave-function, it was shown that only a well defined, discrete spectrum for the cosmological constant is possible. It was also found that among the space-times contributing to wave function, there were two complex conjugate ones that showed a new type of signature change.

It is important to mention that although recent observations point toward a positive cosmological constant, it is still possible that at the very early Universe the cosmological constant be negative. Besides that, we think it is important to understand more about such models which represent bound Universes (analogous to uni-dimensional atoms, in the present situation).

In the present paper, we use the formalism of quantum cosmology in order to quantize three Friedmann-Robertson-Walker models in the presence of a negative cosmological constant and radiation. The radiation is treated by means of the variational formalism developed by Schutz [16]. The models differ from each other by the constant curvature of their spatial sections, which may be positive, negative or zero. They give rise to Wheeler-DeWitt equations for the scale factor, which have the form of the Schrödinger equation for the quartic anharmonic oscillator. We find the eigenvalues and eigenfunctions of those equations by using a method first developed by Chhajlany and Malnev [17]. Then we use the eigenfunctions in order to construct wave packets for each case and evaluate the expectation value of the scale factors as a function of time. In Sec. II we introduce the classical models and solve them analytically, briefly commenting on the general behavior of the classical solutions. In Sec. III we quantize the model by solving the corresponding Wheeler-DeWitt equation. The wave-functional depends on the scale factor a and on the canonical variable associated to the fluid, which in the Schutz variational formalism plays the role of time T . We separate the wave-functional in two parts, one depending solely on the scale factor and the other depending only on the time. The solution in the time sector of the Wheeler-DeWitt equation is trivial, leading to imaginary exponentials of the type $e^{-iE\tau}$, where E is the system energy and $\tau = -T$. The scale factor sector of the Wheeler-DeWitt equation gives rise to the eigenvalue equation for the quartic anharmonic oscillator. We find semi-analytic solutions formed by the product of a decaying exponential term with a polynomial of fixed degree [17]. In Sec. IV we construct wave packets from the eigenfunctions, for each case, and compute the time-dependent, expectation values of the scale factors. We find in all cases that the expectation values of the scale factors show bounded oscillations. Since the expectation values of the scale factors never vanish, we conclude

that these models do not have singularities. We also observe that the energy levels depend on the value of the curvature constant k of the spatial sections. The model with $k < 0$ has the most bounded energy levels, followed by the one with $k = 0$; the model with $k > 0$ have less bounded energy levels. Finally, in Sec. V we summarize the main points and results of our paper.

II. THE CLASSICAL MODELS

Friedmann-Robertson-Walker cosmological models are characterized by the scale factor $a(t)$ and have the following line element,

$$ds^2 = -N(t)^2 dt^2 + a(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right), \quad (1)$$

where $d\Omega^2$ is the line element of the two-dimensional sphere with unitary radius, $N(t)$ is the lapse function and k gives the type of constant curvature of the spatial sections. The curvature is positive for $k = 1$, negative for $k = -1$ and zero for $k = 0$. Here, we are using the natural unit system, where $\hbar = c = G = 1$. The matter content of the model is represented by a perfect fluid with four-velocity $U^\mu = \delta_0^\mu$ in the comoving coordinate system used, plus a negative cosmological constant. The total energy-momentum tensor is given by,

$$T_{\mu,\nu} = (\rho + p)U_\mu U_\nu - pg_{\mu,\nu} - \Lambda g_{\mu,\nu}, \quad (2)$$

where ρ and p are the energy density and pressure of the fluid, respectively. Here, we assume that $p = \rho/3$, which is the equation of the state for radiation. This is justified because the Universe is initially dominated by radiation. Einstein's equations for the metric (1) and the energy momentum tensor (2) are equivalent to the Hamilton equations generated by the super-hamiltonian constraint

$$\mathcal{H} = -\frac{p_a^2}{12} - 3ka^2 + \Lambda a^4 + p_T, \quad (3)$$

where p_a and p_T are the momenta canonically conjugated to a and T the latter being the canonical variable associated to the fluid [11].

The classical dynamics is governed by the Hamilton equations, derived from eq. (3), namely

$$\left\{ \begin{array}{l} \dot{a} = \frac{\partial(N\mathcal{H})}{\partial p_a} = -\frac{N p_a}{6}, \\ \dot{p}_a = -\frac{\partial(N\mathcal{H})}{\partial a} = 6kaN - 4\Lambda a^3 N, \\ \dot{T} = \frac{\partial(N\mathcal{H})}{\partial p_T} = N, \\ \dot{p}_T = -\frac{\partial(N\mathcal{H})}{\partial T} = 0. \end{array} \right. \quad (4)$$

We also have the constraint equation $\mathcal{H} = 0$.

Choosing the gauge $N = 1$, we have the following solutions for the system (4):

$$T(\tau) = \tau + c_1, a(\tau) = \frac{\sqrt{6\beta}}{\sqrt{3k + \sqrt{9k^2 - 12\Lambda\beta}}} sn \left(\frac{\sqrt{18k + 6\sqrt{9k^2 - 12\Lambda\beta}}(\tau - \tau_0)}{6}, \sigma \right), \quad (5)$$

where c_1 , β and τ_0 are integration constants, sn is the Jacobi's elliptic sine [18] of modulus σ given by

$$\sigma = \frac{\sqrt{2}}{2} \sqrt{\frac{-2\beta\Lambda + 3k^2 - k\sqrt{9k^2 - 12\Lambda\beta}}{\Lambda\beta}}. \quad (6)$$

In the case of the models studied here, for which $\Lambda < 0$, Eqs. (5) and (6) imply that the scale factor performs bounded oscillations, for all values of k . When the scale factor vanishes we have the formation of a singularity which may be either a *Big Bang* or a *Big Crunch*. For the sake of completeness, we mention that for $\Lambda = 0$, the case studied in Ref. [6] is recovered.

III. THE QUANTIZATION OF THE MODELS

We wish to quantize the models following the Dirac formalism for quantizing constrained systems [19]. First we introduce a wave-function which is a function of the canonical variables $\hat{a}$ and $\hat{T}$,

$$\Psi = \Psi(\hat{a}, \hat{T}). \quad (7)$$

Then, we impose the appropriate commutators between the operators $\hat{a}$ and $\hat{T}$ and their conjugate momenta $\hat{P}_a$ and $\hat{P}_T$. Working in the Schrödinger picture, the operators $\hat{a}$ and $\hat{T}$ are simply multiplication operators, while their conjugate momenta are represented by the differential operators

$$p_a \rightarrow -i \frac{\partial}{\partial a} \ , \quad p_T \rightarrow -i \frac{\partial}{\partial T} \ . \quad (8)$$

Finally, we demand that $\mathcal{H}$, the super-hamiltonian operator corresponding to (3), annihilate the wave-function Ψ , which leads to Wheeler-DeWitt equation

$$\left(\frac{1}{12} \frac{\partial^2}{\partial a^2} - 3ka^2 + \Lambda a^4 \right) \Psi(a, \tau) = -i \frac{\partial}{\partial \tau} \Psi(a, \tau), \quad (9)$$

where the new variable $\tau = -T$ has been introduced. In order to avoid possible contributions from boundary terms at spatial infinity, we shall consider compact tri-dimensional spatial sections in the cases $k = 0$ and $k = -1$.

The operator $\hat{\mathcal{H}}$ is self-adjoint [6] with respect to the internal product,

$$(\Psi, \Phi) = \int_0^\infty da \ \Psi(a, \tau)^* \Phi(a, \tau), \quad (10)$$

if the wave functions are restricted to the set of those satisfying either $\Psi(0, \tau) = 0$ or $\Psi'(0, \tau) = 0$, where the prime $'$ means the partial derivative with respect to a .

The Wheeler-DeWitt equation (9) is the Schrödinger equation for the quartic anharmonic oscillator and may be solved by writing $\Psi(a, \tau)$ as

$$\Psi(a, \tau) = e^{-iE\tau} \eta(a) \quad (11)$$

where $\eta(a)$ depends solely on a . Then $\eta(a)$ satisfies the eigenvalue equation

$$-\frac{d^2 \eta(a)}{da^2} + V_e(a) \eta(a) = 12E \eta(a), \quad (12)$$

where the effective potential $V_e(a)$ is given by

$$V_e(a) = 36ka^2 - 12\Lambda a^4. \quad (13)$$

A. The Method of Chhajlany and Malnev

The method of Chhajlany and Malnev [17] starts with the addition of an extra term to the original anharmonic oscillator potential, so that the modified Hamiltonian admits a subset of manifestly normalizable solutions. In the case we are considering, the extra term to be added to

the effective potential (13) is proportional to a^6 . In terms of that new enlarged potential, the eigenvalue equation (12) may be re-written as

$$\eta''(a) + (\varepsilon - \alpha a^2 - b a^4 - c a^6) \eta(a) = 0 \quad (14)$$

where $\varepsilon = 12E$, $\alpha = 36k$, $b = -12\Lambda$, c is a parameter to be determined by the method. The *Ansatz* for the solution of Eq. (14) takes the form

$$\eta(a) = N \exp\left(-\frac{c}{4}a^4 - \frac{\gamma}{2}a^2\right) v(a), \quad (15)$$

and has finite norm for $c > 0$. Here, $v(a)$ is a polynomial of a certain degree, yet to be chosen; the parameter γ is to be chosen according to our convenience, as we shall see; N is a normalization factor. The method is based on the fact, shown in Ref. [17], that the larger the degree of the polynomial $v(a)$, the smaller c is. Therefore, if one increases the order of $v(a)$, the energy eigenvalues predicted by the present method tend monotonically, from above, to the energy eigenvalues of the original problem. One important property of the method is that the convergence is very fast. This means that one does not need to use a polynomial of very large order to obtain a good agreement with the energies of anharmonic oscillators already computed in the literature by other methods.

The next step is the substitution of the *Ansatz* (15) into the differential equation (14), which gives rise to the following equation for the polynomial $v(a)$:

$$v''(a) + 2(ca^3 + \gamma a)v'(a) + [\varepsilon - \gamma + (\gamma^2 + \alpha - 3c)a^2]v(a) = 0. \quad (16)$$

Next, writing $v(a) = \sum_n \beta_n a^n$ and inserting it in Eq. (16) along with the condition

$$2c\gamma = b \quad (17)$$

on c and γ , [17] we manage to find:

$$(\varepsilon - \gamma)\beta_0 + 2\beta_2 = 0, \quad (\varepsilon - 3\gamma)\beta_1 + 6\beta_3 = 0, \quad (18)$$

and the general recurrence relation for the polynomial coefficients β_n ,

$$(n+4)(n+3)\beta_{n+4} + [\varepsilon - \gamma(2n+5)]\beta_{n+2} + [\gamma^2 - \alpha - c(2n+3)]\beta_n = 0, \quad (19)$$

for $n \geq 0$. The degree of the polynomial $v(a)$ is fixed to be, say K , by imposing the following conditions in (19),

$$\beta_K \neq 0, \quad \beta_{K+2} = \beta_{K+4} = 0. \quad (20)$$

Due to the nature of the recurrence relation (19), it is clear that by fixing K to be even (odd) the resulting polynomial $v(a)$ will be even (odd). Then, the coefficients β_n , $n = 2, 4, \dots, K$ ($n = 3, 5, \dots, K$), will be determined in terms of β_0 (β_1) by the normalization condition. In the present situation, we restrict our attention to the case of an odd polynomial. It means that, $K = 2m + 1$ for $m = 0, 1, 2, \dots$. This condition is imposed in order that our wave-function vanishes at $a = 0$.

Eqs. (19) and (20) require that the coefficient β_K must vanish; then

$$\gamma^2 = \alpha + c(2K + 3). \quad (21)$$

Combining this with (17), we obtain a cubic algebraic equation in the parameter c ,

$$4c^3(2K + 3) + 4\alpha c^2 - b^2 = 0. \quad (22)$$

The solutions of this equation depend on the known parameters b and K . We must find the real, positive root to this equation so that the *Ansatz* Eq. (15) be normalizable. That real positive root, as proved in Ref. [17], is a monotonically decreasing function of K . Therefore, the greater the polynomial degree, the better the agreement between the energy eigenvalues obtained by this method and the actual energy eigenvalues.

Now, by setting the condition $\beta_{K+2} = 0$ in Eq. (19) we may determine the corresponding energy eigenvalues ε and polynomial coefficients β_n . The $(m + 1)$ allowed energy levels of the anharmonic oscillator are obtained as the roots of the equation $D = 0$, where D is the following $(m + 1) \times (m + 1)$ determinant:

$$\begin{vmatrix} (\varepsilon - 3\gamma) & 6 & 0 & 0 & \dots & \dots & 0 \\ \gamma^2 - \alpha - 5c & (\varepsilon - 7\gamma) & 20 & 0 & \dots & \dots & 0 \\ 0 & \gamma^2 - \alpha - 9c & (\varepsilon - 11\gamma) & 42 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & (\gamma^2 - \alpha - (2K + 3)c) & \varepsilon - (2K + 5)\gamma & (K + 4)(K + 3) \end{vmatrix}. \quad (23)$$

The lowest real, positive root will correspond to the ground state energy level and the excited levels will be given by the sequence of higher real, positive roots. Next, we must substitute these values in the set of Eqs. (19) in order to evaluate the coefficients β_n and obtain the appropriate

polynomial $v_l(a)$; the index $l = 0, 1, 2, \dots, m$ represents the energy level, for each of which we shall have an eigenfunction $\eta_l(a)$ and a wave-function $\Psi_l(a, \tau) = \exp(-iE_l\tau)\eta_l(a)$, according to Eqs. (11) and (15).

We construct a general solution to the Wheeler-DeWitt equation (9) by taking linear combinations of the $\Psi_l(a, \tau)$'s,

$$\Theta(a, \tau) = \sum_{l=0}^m A_l(E) \eta_l(a) e^{-iE_l\tau}, \quad (24)$$

where the coefficients $A_l(E)$ will be fixed later. With those combinations we compute the expected value for the scale factor a , following the *many worlds interpretation* of quantum mechanics [20]. In the present situation, we may write the expected value for the scale factor a is

$$\langle a \rangle (\tau) = \frac{\int_0^\infty a |\Theta(a, \tau)|^2 da}{\int_0^\infty |\Theta(a, \tau)|^2 da}. \quad (25)$$

IV. RESULTS

We shall treat, now, each model separately depending on the constant curvature of the spatial sections. The difference from one model to the other will appear in the value of the parameter α in Eq. (14). For all models we shall use the value of $\Lambda = -0.1$, therefore one has $b = 1.2$ in Eq. (14). Also, we shall fix the polynomial degree to be $K = 45$, for all models. It means that, we shall have 23 energy levels and 23 eigenfunctions $\eta_l(a)$. A precision of at least 15 significant digits had to be used, in order to guarantee the orthogonality of the set of functions $\eta_l(a)$'s Eq. (15). The symbolic system Maple has been used, and the precision of calculations was chosen so that the largest number of energy levels be achieved and the corresponding (approximate) eigenfunctions be sufficiently orthogonal.

A. The model with $k = 1$.

In this model $\alpha = 36$ and the spatial sections are S^3 's. Using the values of K and b , we solved Eq. (22) to find $c = 0.090068960669615962974$. Computations have been performed with 20 significant digits. Now, introducing all these quantities in the determinant D , Eq. (23), we obtain the first 23 energy levels; they are listed in Table I. The first lowest energy levels are in agreement with the ones computed, perturbatively, by Landau for the quartic anharmonic potential, equivalent to the

present case [21]. After that, we substitute these values in the set of Eqs. (19) and compute the coefficients β_n . With these β_n , we write the following $\eta(a, \tau)$, according to Eq. (15),

$$\eta_l(a) = N_l e^{-0.022517240167403990744 a^4 - 3.3307811899866029985 a^2} v_l(a), \quad (26)$$

where

$$v_l(a) = \sum_{i=0}^{22} A_{l,2i+1} a^{2i+1}. \quad (27)$$

The coefficients N_l are normalization coefficients and $i = 0, 1, \dots, m$. The N_l 's and the $A_{l,2i+1}$'s for the present model are listed in the appendix A, Tables II to VII.

Next, we construct the wave-packet $\Theta(a, \tau)$ with the aid of the $\eta_l(a)$, according to Eqs. (24), Eq. (26), and the energy levels in Table I. Finally, using the wave-packet $\Theta(a, \tau)$ we compute the expected value for the scale factor a , Eq. (25). The result is shown in Fig. 1; it can be seen that $\langle a \rangle$ does not vanish, therefore we may say that the quantization of this model removed the singularities it had at the classical level. It is clear from Fig. 1, also, that $\langle a \rangle$ performs bounded oscillations. That means that the spatial sections S^3 's oscillate between finite maximum and minimum radius.

B. The model with $k = 0$

In this model $\alpha = 0$ and the spatial sections are some closed three-dimensional solid with zero curvature, locally isometric to R^3 [22]. Here, like in the previous case, we have used 20 significant digits. Introducing the values of K and b in Eq. (22) we obtain $c = 0.15701453260387612225$. Now, using all these quantities in the determinant D Eq. (23), we obtain the first 23 energy levels. They are shown in Table I. The first lowest energy levels are in agreement with the ones computed, perturbatively, by Landau for the quartic anharmonic potential, equivalent to the present case [21]. After that, we substitute these values in the set of Eqs. (19) and compute the coefficients β_n . With these β_n , we write the following $\eta(a, \tau)$ Eq. (15),

$$\eta_l(a) = N_l e^{-0.039253633150969030562 a^4 - 1.9106511672830600300 a^2} v_l(a), \quad (28)$$

where the $v_l(a)$ have the general expression given in Eq. (27) and the coefficients N_l are normalization coefficients. The N_l 's and the $A_{l,2i+1}$'s for the present model are listed in appendix A, Tables VIII to XIII.

Next, we construct the wave-packet $\Theta(a, \tau)$, with the aid of the $\eta_l(a)$ and the energy levels, according to Eqs. (24) and (28), as well as Table I. Using the wave-packet $\Theta(a, \tau)$ we compute the expected value for the scale factor a , as in Eq. (25). The result is shown in Fig. 2. It can be seen that $\langle a \rangle$ does not assume the value zero; therefore we may say that the quantization of this model removed the singularities it had at the classical level. It is clear, also, from Fig. 2, that $\langle a \rangle$ has bounded oscillations, that is, oscillates between finite maximum and minimum values.

C. The model with $k = -1$

In this model $\alpha = -36$ and the spatial sections are some closed three-dimensional solid with negative constant curvature, locally isometric to H^3 [23]. Here, we have used 15 significant digits. Introducing the values of K and b , we solved Eq. (22) to find $c = 0.410111969406177$. Now, using all these quantities in the determinant D Eq. (23), we obtain the first 23 energy levels; they are listed in Table I. After that, we substitute these values in the set of Eqs. (19) and compute the coefficients β_n , used in

$$\eta_l(a) = N_l e^{-0.102527992350000 a^4 - 0.731507545000000 a^2} v_l(a), \quad (29)$$

where the $v_l(a)$ have the general expression given in Eq. (27) and the N_l 's are normalization coefficients. The N_l 's and the $A_{l,2i+1}$'s for the present model are listed in the appendix A, Tables XIV to XVIII. Due to numerical inconsistencies we have considered, in the present case, 18 $v_l(a)$'s corresponding to the first 18 energy levels.

Next, we construct the wave-packet $\Theta(a, \tau)$ with the aid of the $\eta_l(a)$, according to Eqs. (24) and (29) and the energy levels in Table I. Finally, using the wave-packet $\Theta(a, \tau)$ we compute the expected value for the scale factor a , according to (25). The result is shown in Fig. 3. It can be seen that $\langle a \rangle$ does not assume the value zero. Therefore, we may say that the quantization of this model removed the singularities it had at the classical level. It is also clear that $\langle a \rangle$ performs bounded oscillations.

From Table I, we observe that the energy levels depend on the value of the curvature constant of the spatial sections. The model with negative constant curvature has the most bounded energy levels, then one has the model with zero curvature and finally the model with positive constant curvature has the less bounded energy levels.

Observing Eqs. (26), (28) and (29), we see that the wave-functions $\Psi(a, \tau)$ for all three cases

are exponentially damped as $a \rightarrow \infty$ and behave as powers of a in the limit $a \rightarrow 0$. Following Hawking and Page [24], we may say that this behavior of the $\Psi(a, \tau)$'s makes them *excited states* of wormholes.

Level	$k = 1$	$k = 0$	$k = -1$
E_1	1.5103016760578712464	0.34040279742876372340	-10.5531383749465
E_2	3.5509871014722954423	1.0496018816133576012	-8.85096087898748
E_3	5.6230931685080087038	1.9248782225962522139	-7.22478356024350
E_4	7.7256531719671366439	2.9232145294773461677	-5.68029499265656
E_5	9.8577817762723638293	4.0227383298683634726	-4.22519889374100
E_6	12.018664367453930157	5.2097960735548422725	-2.87075159719689
E_7	14.207548216681935132	6.4748734067709572007	-1.63482529734928
E_8	16.423735430300010131	7.8108774877366064367	-0.536430728059501
E_9	18.666574187417793427	9.2122719218977134548	0.481759230982801
E_{10}	20.935469528987599984	10.674588231181872438	1.58319085808971
E_{11}	23.229800589269486517	12.194126672497696030	2.83303176055370
E_{12}	25.549220854858484858	13.767762525648273494	4.21398303307806
E_{13}	27.892697492531834013	15.392811058629387358	5.70783697327336
E_{14}	30.260978882469751228	17.066942189323770737	7.30286131347368
E_{15}	32.651319599290796010	18.788091721482546958	8.99091730871327
E_{16}	35.066840708164574204	20.554451348143454694	10.7657478496350
E_{17}	37.502643477205645295	22.364363322148552407	12.6225505157144
E_{18}	39.963027226267456788	24.216376790909429384	14.5570952852202
E_{19}	42.443644402959303694	26.109132260627772146	16.5661310939634
E_{20}	44.946892054208187300	28.041419503470901850	18.6464752057244
E_{21}	47.470929938318337116	30.012110489502592151	20.7956557497686
E_{22}	50.01608754545613080	32.020172678051097447	23.0112449804346
E_{23}	52.581852999800707709	34.064648529603099090	25.2911792258758

TABLE I: The lowest calculated energy levels for the cases $k = 0$, $k = 1$, and $k = -1$ (in all cases, $\Lambda = -0.1$).

V. CONCLUSIONS.

In the present paper, the formalism of quantum cosmology was employed to quantize three Friedmann-Robertson-Walker models in the presence of a negative cosmological constant and radiation. The variational formalism of Schutz [16] allowed us to ascribe dynamical degrees of freedom to the radiation fluid. The models differ from each other by the constant curvature of the spatial sections, which may be positive, negative or zero. The quantization of the models gave rise to Wheeler-DeWitt equations, for the scale factor, which had the form of the Schrödinger equation for the quartic anharmonic-oscillator. We found the approximate eigenvalues and eigenfunctions of those equations by using a method first developed by Chhajlany and Malnev [17]. After that,

we used the eigenfunctions in order to construct wave-packets for each case and evaluate the time dependent, expected value of the scale factors. We found for all of them that the expected values of the scale factors evolve with bounded oscillations. Since the expectation value of the scale factors never vanish, we concluded that these models do not have singularities. We also observed that the energy levels depend on the value of the curvature constant of the spatial sections. The model with negative curvature constant has the most bounded energy levels, whereas the model with positive constant curvature has the less bounded energy levels.

Acknowledgments

E. V. Corrêa Silva thanks CNPq for partial financial support.

-
- [1] B. S. DeWitt, Phys. Rev. D **160**, 1113 (1967).
 - [2] J. B. Hartle and S. W. Hawking, Phys. Rev. **D28**, 2960 (1983).
 - [3] G. Oliveira-Neto, Phys. Rev. D **58**, 024010 (1998);
 - [4] Y. Fujiwara et al., Class. Quantum Grav. **7** 163 (1992); Phys. Rev. D **44**, 1756 (1991); J. Louko and P. J. Ruback, Class. Quantum Grav. **8**, 91 (1991); J. J. Halliwell and J. Louko, Phys. Rev. D **42**, 3997 (1990);
 - [5] Mariam Bouhmadi-Lopez and Paulo Vargas Moniz, Phys. Rev. D **71**, 063521 (2005); I. G. Moss and W. A. Wright, Phys. Rev. D **29**, 1067 (1984); M. J. Gotay and J. Demaret, Phys. Rev. D **28**, 2402 (1983);
 - [6] N. A. Lemos, J. Math. Phys. **37**, 1449 (1996);
 - [7] A. Vilenkin, Phys. Rev. D **33**, 3560 (1986); **37**, 888 (1988).
 - [8] D. Bohm and B. J. Hiley, *The undivided Universe: an ontological interpretation of quantum theory* (Routledge, London, 1993); P. R. Holland, *The quantum theory of motion: an account of the de Broglie-Bohm interpretation of quantum mechanics* (Cambridge University Press, Cambridge, 1993).
 - [9] C. W. Misner, in *Magic without Magic: John Archibald Wheeler, a Collection of Essays in Honor of his 60th Birthday*, ed. J. Klauder (W. H. Freeman, San Francisco, 1972), 441.
 - [10] N. A. Lemos, G. A. Monerat, Gen. Rel. Grav. **35**, 423 (2003); J. Acacio de Barros and N. Pinto-Neto, Int. J. Mod. Phys. D **7**, 201 (1998); J. Acacio de Barros, N. Pinto-Neto and M. A. Sagiore-Leal, Phys. Lett. A **241**, 229 (1998).
 - [11] F. G. Alvarenga, J. C. Fabris, N. A. Lemos, G. A. Monerat, Gen. Rel. Grav. **34**, 651 (2002).
 - [12] M. Bojowald, Class. Quantum Grav. **20**, 2595 (2003); **19**, 2717 (2002); Phys. Rev. Lett. **87**, 121301 (2001); **86**, 5227 (2001).

- [13] For a recent review see M. Bojowald, *Pramana* **63**, 765 (2004).
- [14] S. Carlip, *Phys. Rev. Lett.* **79** 4071 (1997).
- [15] M. Anderson, S. Carlip, J.G. Ratcliffe, S. Surya and S.T. Tschantz, *Class. Quant. Grav.* **21** 729, (2004).
- [16] Schutz, B. F., *Phys. Rev. D* **2**, 2762, (1970); Schutz, B. F., *Phys. Rev. D* **4**, 3359, (1971).
- [17] S. C. Chhajlany and V. N. Malnev, *Phys. Rev. A*, vol 42, No. 5, 3111 p., (1990); S. C. Chhajlany, D. A. Letov and V. N. Malnev, *J. Phys. A: Math. Gen.* **24**, 2731 (1991).
- [18] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions*, (Dover Publications Inc., New York, 1965), p. 1046.
- [19] P. A. M. Dirac, *Can. J. Math.* **2**, 129 (1950); *Proc. Roy. Soc. London A* **249**, 326 and 333 (1958); *Phys. Rev.* **114**, 924 (1959).
- [20] H. Everett, III, *Rev. Mod. Phys.* **29**, 454 (1957).
- [21] E. M. Lifshitz, L. D. Landau, *Quantum Mechanics: Non-Relativistic Theory, Volume 3, Third Edition (Quantum Mechanics)*, (Butterworth-Heinemann, Oxford, 2003).
- [22] J. A. Wolf, *Spaces of Constant Curvature*, (McGraw-Hill, New York, 1967).
- [23] W. P. Thurston, *The Geometry and Topology of Three-Manifolds*, (Princeton University Press, Princeton, 1982).
- [24] S. W. Hawking and D. N. Page, *Phys. Rev. D* **42**, 2655 (1990).

APPENDIX A: COEFFICIENTS OF THE POLYNOMIALS $v_i(a)$

The following Tables contain the coefficients $A_{l,2i+1}$ introduced in eq. (27), for each case considered. Tables II to VII refer to the case $\Lambda = -0.1$, $k = 1$; Tables VIII to XIII refer to the case $\Lambda = -0.1$, $k = 0$; Tables XIV to XVIII refer to the case $\Lambda = -0.1$, $k = -1$. The normalization coefficients N_l can be found in the respective captions.

i	Coefficients $A_{1,2i+1}$	Coefficients $A_{2,2i+1}$	Coefficients $A_{3,2i+1}$	Coefficients $A_{4,2i+1}$
0	1	1	1	1
1	0.31017783787086050567	-3.7711930129579878862	-7.9154051470294144092	-12.120525153947670289
2	0.04581346355795285829	-1.1541419051991141135	7.853995135600127347	27.527508697085798433
3	0.004286539733354114758	-0.1633354878078236419	2.510476703458296824	-10.551810905706899599
4	0.0002851177209318121384	-0.014511494987299668478	0.3453281989577501333	-3.812327078892842085
5	0.000014340884997309905358	-0.0009116635877726964358	0.02928647013334059913	-0.5169340736618381480
6	0.000000566441403839108	-0.0000431229586177813	0.001740679187283938	-0.042024926844112081
7	0.000000018009305087394	-0.0000015951640298011	0.000077408641677853	-0.002364735333609904
8	0.000000000468665873116	-0.0000000472918331013	0.000002677318849025	-0.000098747186611166
9	1.0096366866 $\times 10^{-11}$	-0.0000000114220503333	0.0000000738149538296	-0.000003185308801166
10	1.813735938 $\times 10^{-13}$	-2.271666337 $\times 10^{-11}$	0.0000000016484458275	-0.000000081371293283
11	2.7264391 $\times 10^{-15}$	-3.7452535 $\times 10^{-13}$	3.01217556 $\times 10^{-11}$	-0.000000001672240001
12	3.33586 $\times 10^{-17}$	-5.15166 $\times 10^{-15}$	4.527534 $\times 10^{-13}$	-2.79071173 $\times 10^{-11}$
13	-4.521 $\times 10^{-20}$	-6.6200 $\times 10^{-17}$	5.54736 $\times 10^{-15}$	-3.800602 $\times 10^{-13}$
14	-1.7706 $\times 10^{-19}$	-3.510 $\times 10^{-18}$	3.324 $\times 10^{-17}$	-4.2752 $\times 10^{-15}$
15	-7.060 $\times 10^{-20}$	-1.0944 $\times 10^{-18}$	-7.71 $\times 10^{-18}$	-5.552 $\times 10^{-17}$
16	-2.5166326155344480568 $\times 10^{-20}$	-3.8103863313805363486 $\times 10^{-19}$	-2.6663403598954915102 $\times 10^{-18}$	-5.4958151805952792446 $\times 10^{-18}$
17	-8.2632648107601721899 $\times 10^{-21}$	-1.2709238053250021166 $\times 10^{-19}$	-8.3203099235009615604 $\times 10^{-19}$	-1.5357192261084095937 $\times 10^{-18}$
18	-2.8853679863959393083 $\times 10^{-21}$	-4.1339650865318377254 $\times 10^{-20}$	-2.4995310291072082197 $\times 10^{-19}$	-4.2585576822142560559 $\times 10^{-19}$
19	-9.7000912539108576359 $\times 10^{-22}$	-1.2385504232519897984 $\times 10^{-20}$	-7.1369867141313118842 $\times 10^{-20}$	-1.1454286664270462490 $\times 10^{-19}$
20	-2.8079252483628921590 $\times 10^{-22}$	-3.5730297879647888446 $\times 10^{-21}$	-1.9694920031159536822 $\times 10^{-20}$	-2.9769568394984189501 $\times 10^{-20}$
21	-8.8765902812953993115 $\times 10^{-23}$	-1.0510940838473940756 $\times 10^{-21}$	-5.3288366402638849473 $\times 10^{-21}$	-7.687451297285508365 $\times 10^{-21}$
22	-2.3906161697064305868 $\times 10^{-23}$	-2.8689415174657205262 $\times 10^{-22}$	-1.3846743125110228663 $\times 10^{-21}$	-1.8976098904628921420 $\times 10^{-21}$

TABLE II: The case $\Lambda = -0.1$, $k = 1$: coefficients $A_{1,2i+1}$, $A_{2,2i+1}$, $A_{3,2i+1}$ and $A_{4,2i+1}$ of the polinomials v_1 , v_2 , v_3 and v_4 . The coefficients of the normalization are $N_1 = 5.8034051393293824531$, $N_2 = 7.1778933987585237889$, $N_3 = 8.1009761694435305313$ and $N_4 = 8.8293734529556238792$.

i	Coefficients $A_{5,2i+1}$	Coefficients $A_{6,2i+1}$	Coefficients $A_{7,2i+1}$	Coefficients $A_{8,2i+1}$
0	1	1	1	1
1	-16.384782362558124660	-20.706547544921257315	-25.084315243377267265	-29.516689670613417263
2	58.312374535479288149	100.64443831320425508	154.95041179125834137	221.64873340822231830
3	-59.54850867092679165	-166.27947628665417837	-354.13029662189362099	-648.05926695036679593
4	9.35789837391042752	92.23715065066104101	331.57590893769938406	852.35062705243081723
5	4.409713597601337702	-4.45598233475616519	-110.35457194930656034	-506.0681988766809666
6	0.59818020215651014	-4.09117137488257791	-1.56360443105183866	106.8793989497608568
7	0.04676720567471725	-0.56433583205511656	3.14873326051023910	5.9941189879547596
8	0.00249049073418037	-0.04252260991596205	0.45008779793574406	-2.05739270958767402
9	0.000097440326882724	-0.00213981700876802	0.03271392094015454	-0.31170440703586862
10	0.000002921073818425	-0.00007817947147838	0.00155121718457568	-0.02183733966183077
11	0.000000068813279087	-0.000002167823114800	0.00005267868723725	-0.00097142641061071
12	0.000000001293537112	-0.000000046808363285	0.000001342896942912	-0.00003047372169922
13	1.95721139 $\times 10^{-11}$	-0.000000000798594075	0.000000026377008433	-0.000000708587224832
14	2.406330 $\times 10^{-13}$	-1.08103419 $\times 10^{-11}$	0.000000000405172515	-0.000000012530964318
15	2.8483 $\times 10^{-15}$	-1.078220 $\times 10^{-13}$	4.9900537 $\times 10^{-12}$	-0.0000000001697070756
16	1.5617272916640256937 $\times 10^{-16}$	1.380268516222927925 $\times 10^{-15}$	7.165706651611258944 $\times 10^{-14}$	-1.570909246541791477 $\times 10^{-12}$
17	3.699521211534707235 $\times 10^{-17}$	5.789009553323564153 $\times 10^{-16}$	6.03431426543476250 $\times 10^{-15}$	3.0635698429937474 $\times 10^{-14}$
18	9.608892062179437947 $\times 10^{-18}$	1.406502616814589012 $\times 10^{-16}$	1.25407781422107830 $\times 10^{-15}$	8.8947211205413485 $\times 10^{-15}$
19	2.427534289107907126 $\times 10^{-18}$	3.302474693051297305 $\times 10^{-17}$	2.71172197801927290 $\times 10^{-16}$	1.7785141700571135 $\times 10^{-15}$
20	5.955763963500422277 $\times 10^{-19}$	7.57138310970926381 $\times 10^{-18}$	5.7727046580805942 $\times 10^{-17}$	3.4916244211180328 $\times 10^{-16}$
21	1.415564509865121219 $\times 10^{-19}$	1.692656131320643543 $\times 10^{-18}$	1.20592188277216537 $\times 10^{-17}$	6.772423485917385 $\times 10^{-17}$
22	3.2584165617389642358 $\times 10^{-20}$	3.6905326034954039964 $\times 10^{-19}$	2.4692233577815985073 $\times 10^{-18}$	1.2953462920958676956 $\times 10^{-17}$

TABLE III: The case $\Lambda = -0.1$, $k = 1$: coefficients $A_{5,2i+1}$, $A_{6,2i+1}$, $A_{7,2i+1}$ and $A_{8,2i+1}$ of the polinomials v_5 , v_6 , v_7 and v_8 . The coefficients of the normalization are $N_5 = 9.4465594769581691233$, $N_6 = 9.9907608238646431121$, $N_7 = 10.482871871428757467$ and $N_8 = 10.935645654327260289$.

i	Coefficients $A_{9,2i+1}$	Coefficients $A_{10,2i+1}$	Coefficients $A_{11,2i+1}$	Coefficients $A_{12,2i+1}$
0	1	1	1	1
1	-34.002367184848983855	-38.540157867988596970	-43.128819988552370035	-47.767660519730366718
2	301.15021081131710573	393.85929396557445690	500.17116570886464547	620.48706397200240588
3	-1074.5861450716873336	-1661.7872217348979949	-2439.2637603904004098	-3438.2520498616334939
4	1821.6641872613443414	3452.6958410202546019	6008.6072441957345911	9806.9867235942380195
5	-1546.6135676876763965	-3809.3067552029814704	-8161.2736489861159871	-15835.705001515897835
6	626.0285291644673913	2249.7333239006545920	6318.0405997660246921	15161.530656034483683
7	-86.4681405913189308	-653.4186254833888626	-2737.3334802679376352	-8687.6607088159865421
8	-7.7070589894374189	59.7450684776875324	593.1995682188562094	2875.3051765536900470
9	1.16000437460191013	7.1681695347971782	-35.8441748442604442	-480.1213500044692684
10	0.19143777716358755	-0.57113494102661983	-5.5060245096840928	18.9573818265190906
11	0.01288929254728267	-0.10610473298349592	0.24810332800424582	3.68937193444829489
12	0.00053453455552767	-0.00682545444872026	0.05387180639994624	-0.09661243436302107
13	0.000015365732576736	-0.00026161239759925	0.00327916311924026	-0.02538990883522057
14	0.00000322761471282	-0.000006818839602350	0.000114852787218929	-0.001441041209603695
15	0.000000005090659309	-0.000000127615536791	0.000002679703174836	-0.000045418784915808
16	$6.23793384554829457 \times 10^{-11}$	-0.000000001746106498202539696	0.0000000412547238772015166	-0.000000930805731905662193
17	$8.6575190851395808 \times 10^{-13}$	$-1.5820289883099125 \times 10^{-11}$	0.00000000053441175665178327	-0.0000000130438640811481176
18	$6.058979416709158 \times 10^{-14}$	$2.000038405818014 \times 10^{-13}$	$6.310684727048987 \times 10^{-12}$	-0.0000000001187393590485689
19	$1.0215990409800383 \times 10^{-14}$	$5.27514494520875 \times 10^{-14}$	$2.90888564322591 \times 10^{-13}$	$3.8940582890413 \times 10^{-13}$
20	$1.829640531826615 \times 10^{-15}$	$8.67098043007747 \times 10^{-15}$	$3.83550545914211 \times 10^{-14}$	$1.5649846755718 \times 10^{-13}$
21	$3.270932265719669 \times 10^{-16}$	$1.41685618892377 \times 10^{-15}$	$5.6481920779378 \times 10^{-15}$	$2.1111601530203 \times 10^{-14}$
22	$5.8067651622857174343 \times 10^{-17}$	$2.3178747497526342364 \times 10^{-16}$	$8.4410012164581453358 \times 10^{-16}$	$2.8535038929738800685 \times 10^{-15}$

TABLE IV: The case $\Lambda = -0.1$, $k = 1$: coefficients $A_{9,2i+1}$, $A_{10,2i+1}$, $A_{11,2i+1}$ and $A_{12,2i+1}$ of the polynomials v_9 , v_{10} , v_{11} and v_{12} . The coefficients of the normalization are $N_9 = 11.357473916077508835$, $N_{10} = 11.754220307373787325$, $N_{11} = 12.130088843440861937$ and $N_{12} = 12.488270658573768104$.

i	Coefficients $A_{13,2i+1}$	Coefficients $A_{14,2i+1}$	Coefficients $A_{15,2i+1}$	Coefficients $A_{16,2i+1}$
0	1	1	1	1
1	-52.454613795077065028	-57.191176574952899458	-61.971858008594989022	-66.802900226342545410
2	755.16371270759864284	904.65638191366384671	1069.1911724988718290	1249.3896027640753475
3	-4691.1786486789463820	-6233.0019360858040813	-8097.8534617243312174	-10325.918637224436785
4	15222.159586897177725	22695.269368957207490	32722.191703657035811	45893.949491372741828
5	-28511.535588496471157	-48419.757316339959116	-78400.032812948401888	-122116.39684597553325
6	32538.633947802319935	64210.863046864482262	118551.52404606112225	207557.92486531471539
7	-23161.853380843369227	-54625.432365110668228	-117367.86905261772511	-234576.48356569840002
8	10227.480262043182441	30093.669911892066791	77447.628014036355538	180411.03777136843423
9	-2671.0800924104234944	-10573.928314703002037	-34075.022670025208808	-95280.641820767751010
10	353.5915507866116550	2240.0578702285590691	9780.4182354247769263	34357.916918381656131
11	-9.0038128802620698	-241.6086156778874408	-1721.2573047719675453	-8240.9083549293767413
12	-2.22338842366103368	4.00870518474146025	155.36288454585560845	1231.7953979185424375
13	0.03490705538413532	1.23038651087727620	-1.80811935476349017	-95.71694844973517166
14	0.01122655434357897	-0.01285528560006987	-0.63270275783899063	0.94912229292304996
15	0.000582306454656128	-0.004703102634057810	0.005476570471229182	0.30538010701217688
16	0.00001621120697813077765	-0.0002167549854023615158	0.0018713746378842268557	-0.00291544473702381013
17	0.000000286414799028962413	-0.00000518620428293573924	0.00007411339194162508172	-0.000709711959014798152
18	0.000000003420615524128955	-0.00000007590009610821422	0.00000147853815770511447	-0.000023065063153326174
19	$3.24792731721724 \times 10^{-11}$	-0.000000000686192176752252	0.000000017464288701285557	-0.0000003639775018973054
20	$7.877917012049 \times 10^{-13}$	$-1.560853760088 \times 10^{-12}$	0.000000000132503203659754	-0.0000000031380577144596
21	$7.574158403458 \times 10^{-14}$	$2.466441247840 \times 10^{-13}$	$1.3674223184521 \times 10^{-12}$	$-1.171890220165 \times 10^{-11}$
22	$9.079369472488977225 \times 10^{-15}$	$2.7527477962670556412 \times 10^{-14}$	$8.143665471068961802 \times 10^{-14}$	$2.023798389867859016 \times 10^{-13}$

TABLE V: The case $\Lambda = -0.1$, $k = 1$: coefficients $A_{13,2i+1}$, $A_{14,2i+1}$, $A_{15,2i+1}$ and $A_{16,2i+1}$ of the polynomials v_{13} , v_{14} , v_{15} and v_{16} . The coefficients of the normalization are $N_{13} = 12.831657346516500646$, $N_{14} = 13.160199699505029703$, $N_{15} = 13.481183098459831301$ and $N_{16} = 13.783901938149622931$.

i	Coefficientes $A_{17,2i+1}$	Coefficientes $A_{18,2i+1}$	Coefficientes $A_{19,2i+1}$	Coefficientes $A_{20,2i+1}$
0	1	1	1	1
1	-71.674505764424687592	-76.595273262548310578	-81.556507615932004390	-86.563002918429771602
2	1445.2812913047251820	1657.6056242524762461	1886.3841243300680194	2132.2207701020935152
3	-12951.762075898028761	-16020.740104978502468	-19569.862939305032690	-23646.245366475767022
4	62834.794759969799382	84306.333013897444312	111087.71291195023833	144107.29200810027984
5	-183963.75221484100112	-269539.48654730779802	-385321.01208259817624	-539314.72674103090793
6	347409.38740624855152	560351.91063946410830	874970.62323112283525	1328975.0572268366759
7	-441584.34608490955678	-791825.42722176937838	-1361472.4047563478949	-2259352.8071906066129
8	387990.66363169029593	783000.60429136018579	1497172.7240646876630	2736670.8438980238668
9	-239063.98678343357541	-551618.42718575789884	-1187471.2980357767532	-2414563.4814397670821
10	103619.90361692807112	279218.51883627417348	687499.12123336995985	1574324.4179742275843
11	-31286.43769518295557	-101454.73705241262000	-291829.66149555115473	-764549.93448862474893
12	6393.8838281755775839	26135.501634440661231	90489.250677174203654	276979.77502284607572
13	-827.92773276506180597	-4632.6347050968838681	-20212.254691993688141	-74446.727185366485745
14	56.91755063260658843	529.70099076132116387	3156.5078170609347092	14630.127421681580902
15	-0.60186346671718578	-33.12977276170413576	-324.23366713806025380	-2042.4107514564805240
16	-0.13804887378499444466	0.432796730259634788	18.90939127874898872	191.55056588041964646
17	0.00170555532562603351	0.0581247473296715877	-0.313931804144568406	-10.64761982516368418
18	0.000253407010392459341	-0.0009897386034172207	-0.022119579552745663	0.21985595919567848
19	0.000006405499376370944	-0.00008404838802303226	0.0005264609495686830	0.007045958332765000
20	0.0000000750710062604530	-0.00000152398837574875	0.0000248748101605601	-0.000249470805616634
21	0.0000000004429189405509	-0.000000011599026864238	0.00000028525947585523	-0.0000060880619974117
22	$1.65463769026912201 \times 10^{-12}$	$-3.12048788790252230 \times 10^{-11}$	0.00000001066066931450985327	-0.0000000327991012433832455

TABLE VI: The case $\Lambda = -0.1$, $k = 1$: coefficients $A_{17,2i+1}$, $A_{18,2i+1}$, $A_{19,2i+1}$ and $A_{20,2i+1}$ of the polynomials v_{17} , v_{18} , v_{19} and v_{20} . The coefficients of the normalization are $N_{17} = 14.088838258318067966$, $N_{18} = 14.371929657653422158$, $N_{19} = 14.659291915176228163$ and $N_{20} = 14.930156447596541712$.

i	Coefficients $A_{21,2i+1}$	Coefficients $A_{22,2i+1}$	Coefficients $A_{23,2i+1}$
0	1	1	1
1	-71.674505764424687592	-76.595273262548310578	-81.556507615932004390
2	1445.2812913047251820	1657.6056242524762461	1886.3841243300680194
3	-12951.762075898028761	-16020.740104978502468	-19569.862939305032690
4	62834.794759969799382	84306.333013897444312	111087.71291195023833
5	-183963.75221484100112	-269539.48654730779802	-385321.01208259817624
6	347409.38740624855152	560351.91063946410830	874970.62323112283525
7	-441584.34608490955678	-791825.42722176937838	-1361472.4047563478949
8	387990.66363169029593	783000.60429136018579	1497172.7240646876630
9	-239063.98678343357541	-551618.42718575789884	-1187471.2980357767532
10	103619.90361692807112	279218.51883627417348	687499.12123336995985
11	-31286.43769518295557	-101454.73705241262000	-291829.66149555115473
12	6393.8838281755775839	26135.501634440661231	90489.250677174203654
13	-827.92773276506180597	-4632.6347050968838681	-20212.254691993688141
14	56.91755063260658843	529.70099076132116387	3156.5078170609347092
15	-0.60186346671718578	-33.12977276170413576	-324.23366713806025380
16	-0.13804887378499444466	0.432796730259634788	18.90939127874898872
17	0.00170555532562603351	0.0581247473296715877	-0.313931804144568406
18	0.000253407010392459341	-0.0009897386034172207	-0.022119579552745663
19	0.000006405499376370944	-0.00008404838802303226	0.0005264609495686830
20	0.0000000750710062604530	-0.00000152398837574875	0.0000248748101605601
21	0.0000000004429189405509	-0.000000011599026864238	0.00000028525947585523
22	$1.65463769026912201 \times 10^{-12}$	$-3.12048788790252230 \times 10^{-11}$	0.00000001066066931450985327

TABLE VII: The case $\Lambda = -0.1$, $k = 1$: coefficients $A_{21,2i+1}$, $A_{22,2i+1}$ and $A_{23,2i+1}$ of the polynomials v_{21} , v_{22} and v_{23} . The coefficients of the normalization are $N_{21} = 15.196350124389910569$, $N_{22} = 15.443515484903478896$ and $N_{23} = 15.645567647056785661$.

i	Coefficients $A_{1,2i+1}$	Coefficients $A_{2,2i+1}$	Coefficients $A_{3,2i+1}$	Coefficients $A_{4,2i+1}$
0	1	1	1	1
1	1.2298455724255325832	-0.18855259594365517233	-1.9391052779094443978	-3.9357778916716323053
2	0.70281444756044962535	-0.8243016140395259213	-1.0448066649011448432	0.9482808911585986722
3	0.24882730009488482050	-0.51856758078952620652	0.1378821684225774495	1.3929964156767481609
4	0.061362228801690876509	-0.17831099849711656507	0.24781182769474065702	0.26486205576426314590
5	0.011229566129419549313	-0.041020332973195298495	0.096571463561412127319	-0.06075921121492778053
6	0.0015858480631393467966	-0.006876928456515549490	0.022150497291839980024	-0.039763176718287530660
7	0.00017735362766665490129	-0.0008806578965821418589	0.0035364032024917661879	-0.0098048157493233599636
8	0.000015988383854415975967	-0.00008869561348138795822	0.0004215040272918221427	-0.0015366389100990166141
9	0.00000117614033597329686	-0.0000071606437006450870	0.0000389537207381306872	-0.0001732328915718124902
10	0.00000007117596314205952	-0.00000046924399378806075	0.0000028555317909651724	-0.0000148306955832690722
11	0.000000003560689954269945	-0.00000002515643731461728	0.00000016838604370235678	-0.00000099264167067812283
12	0.0000000001475461488138725	-0.000000001107875969030644	0.000000008052643519724566	-0.00000005280787155980257
13	$5.0605642376533 \times 10^{-12}$	$-4.01129360183140 \times 10^{-11}$	$0.000000000313430341951797$	$-0.000000002252576562969348$
14	$1.4315256162425 \times 10^{-13}$	$-1.1912125948781 \times 10^{-12}$	$9.9250001786209 \times 10^{-12}$	$-7.72738658315285 \times 10^{-11}$
15	$3.31721811120 \times 10^{-15}$	$-2.884315756394 \times 10^{-14}$	$2.5455256253983 \times 10^{-13}$	$-2.1274442573091 \times 10^{-12}$
16	$6.2290296428 \times 10^{-17}$	$-5.63701481638 \times 10^{-16}$	$5.24043213312 \times 10^{-15}$	$-4.666452306035 \times 10^{-14}$
17	$9.325441740 \times 10^{-19}$	$-8.753563734 \times 10^{-18}$	$8.5319740794 \times 10^{-17}$	$-8.04499352583 \times 10^{-16}$
18	$1.086336997 \times 10^{-20}$	$-1.054610245 \times 10^{-19}$	$1.0734023077 \times 10^{-18}$	$-1.06620225448 \times 10^{-17}$
19	9.485949×10^{-23}	$-9.4996313 \times 10^{-22}$	$1.00619408 \times 10^{-20}$	$-1.048181689 \times 10^{-19}$
20	5.83919×10^{-25}	$-6.018252 \times 10^{-24}$	6.614357×10^{-23}	$-7.1979226 \times 10^{-22}$
21	2.2587×10^{-27}	-2.3864×10^{-26}	2.72624×10^{-25}	$-3.073159 \times 10^{-24}$
22	4.85×10^{-30}	-3.79×10^{-29}	6.312×10^{-28}	-4.9228×10^{-27}

TABLE VIII: The case $\Lambda = -0.1$, $k = 0$: coefficients $A_{1,2i+1}$, $A_{2,2i+1}$, $A_{3,2i+1}$ and $A_{4,2i+1}$ of the polinomys v_1 , v_2 , v_3 and v_4 . The coefficients of the normalization are $N_1 = 2.1407112737145309126$, $N_2 = 3.2975672670529412541$, $N_3 = 4.1572580515879983879$ and $N_4 = 4.8776996852430886717$.

i	Coefficients $A_{5,2i+1}$	Coefficients $A_{6,2i+1}$	Coefficients $A_{7,2i+1}$	Coefficients $A_{8,2i+1}$
0	1	1	1	1
1	-6.1348254924536669152	-8.5089409798266245150	-11.039095646258854371	-13.711103808190152843
2	5.911356608106969186	14.526711829815446250	27.430881469808772742	45.228591749889808489
3	1.048461113886525862	-4.412562811395891946	-19.826241542993934663	-51.364469972481314193
4	-0.8995614524956676092	-2.215775126916415019	0.826029728265588547	18.084829935024654405
5	-0.31272201167583072358	0.2754965915900852529	2.112554853993957234	2.098921351752975453
6	-0.01422790600763145292	0.20538109262592843443	0.07815218589511349951	-1.3891586577302272466
7	0.012179925838702809322	0.025756335876411061922	-0.09792697564027372845	-0.16919167853767788552
8	0.0036685806203861655213	-0.002290361001058895760	-0.017562627232355277223	0.03593924509220542531
9	0.000581339324052616344	-0.00118650597006258730	-0.00018063403724606157	0.00886562542581478584
10	0.000062660478934633456	-0.000196451799277369343	0.000336999686606672515	0.00041491502516795559
11	0.0000049897731210930381	-0.000020377384978350386	0.000060602588364133200	-0.000085175283410706295
12	0.00000030507598448885760	-0.0000015087682548176592	0.0000060595073530437073	-0.000017401914257678935
13	0.00000001460930559723988	-0.00000008395147832486714	0.0000004151278365955338	-0.0000016693173630631525
14	0.000000000553170023976602	-0.00000000359585742236137	0.00000002085721988055981	-0.00000010478203231223269
15	$1.6595082577133 \times 10^{-11}$	$-0.000000000119751354995018$	$0.000000000790654647053799$	$-0.00000000469537115086084$
16	$3.926691338187 \times 10^{-13}$	$-3.1016711682977 \times 10^{-12}$	$2.2835966945664 \times 10^{-11}$	$-0.000000000155113615814839$
17	$7.24370011893 \times 10^{-15}$	$-6.19561204242 \times 10^{-14}$	$5.010676197816 \times 10^{-13}$	$-3.8081259712585 \times 10^{-12}$
18	$1.02040365374 \times 10^{-16}$	$-9.3689489119 \times 10^{-16}$	$8.22808582580 \times 10^{-15}$	$-6.88680006219 \times 10^{-14}$
19	$1.060342406 \times 10^{-18}$	$-1.0377392653 \times 10^{-17}$	$9.807188694 \times 10^{-17}$	$-8.9322016791 \times 10^{-16}$
20	$7.6611823 \times 10^{-21}$	$-7.94452548 \times 10^{-20}$	$8.02026022 \times 10^{-19}$	$-7.874055026 \times 10^{-18}$
21	3.443267×10^{-23}	$-3.753929 \times 10^{-22}$	$4.0289718 \times 10^{-21}$	$-4.22907569 \times 10^{-20}$
22	8.1693×10^{-26}	-7.7752×10^{-25}	9.57681×10^{-24}	$-1.035546 \times 10^{-22}$

TABLE IX: The case $\Lambda = -0.1$, $k = 0$: coefficients $A_{5,2i+1}$, $A_{6,2i+1}$, $A_{7,2i+1}$ and $A_{8,2i+1}$ of the polinomys v_5 , v_6 , v_7 and v_8 . The coefficients of the normalization are $N_5 = 5.5127641866644550323$, $N_6 = 6.0885073250502348952$, $N_7 = 6.6199175722251377252$ and $N_8 = 7.1165584094285250131$.

i	Coefficients $A_{9,2i+1}$	Coefficients $A_{10,2i+1}$	Coefficients $A_{11,2i+1}$	Coefficients $A_{12,2i+1}$
0	1	1	1	1
1	-16.513892676512366880	-19.438525295080684847	-22.477602177712332030	-25.624873884013486958
2	68.500816128447368264	97.809919332667553061	133.70317321896426872	176.71530644883325203
3	-106.55682738798604833	-194.31462763518727966	-324.95562820472674522	-510.22884819063511319
4	66.82399915663488053	173.94601726151463769	378.40039317842692369	733.75951884021536989
5	-11.49015530870562177	-66.66856760315107297	-218.36140595295844410	-562.39713852806645723
6	-3.173797462030778022	4.57659601834016473	54.37892239826436592	225.58435351580697565
7	0.6956085962224014391	2.846529595819562843	-0.07259654297061238	-38.04805538744838408
8	0.13749152817338703835	-0.2699092404950981048	-2.001569064665355247	-1.794224678997464712
9	-0.0098352843123713145	-0.0828711087596463085	0.0756477908454371566	1.2131746963605570921
10	-0.00377960440222442262	0.00151974846501950219	0.04238992696925214174	-0.0091803457551753205
11	-0.000242928582381203110	0.00144598631945890593	0.00028616572311268950	-0.01955936534615395811
12	0.000019517596540780945	0.000105038841810993914	-0.000516598070399832600	-0.000353719702095261487
13	0.0000047356176686937723	-0.0000042366253287158835	-0.000038549025817450602	0.000177539291967760406
14	0.0000004300505576264660	-0.0000012406806155959293	0.0000009648831285926503	0.0000125658395523811823
15	0.00000002435348177638940	-0.00000010413682689576750	0.00000031639556514865196	-0.0000002650886578360362
16	0.000000000956365820294936	-0.000000005205438956721505	0.00000002369666978144363	-0.00000007874855629121539
17	$2.6944771045870 \times 10^{-11}$	-0.000000000174886873756666	0.000000001015518810192359	-0.000000005030287511806847
18	$5.464426457482 \times 10^{-13}$	$-4.0772785430683 \times 10^{-12}$	$2.82471588404887 \times 10^{-11}$	-0.0000000001778964228815842
19	$7.81851571985 \times 10^{-15}$	$-6.549507537492 \times 10^{-14}$	$5.216954308548 \times 10^{-13}$	$-3.9118136989119 \times 10^{-12}$
20	$7.5103400955 \times 10^{-17}$	$-6.9461889320 \times 10^{-16}$	$6.21222460816 \times 10^{-15}$	$-5.348131653543 \times 10^{-14}$
21	$4.35481724 \times 10^{-19}$	$-4.390674715 \times 10^{-18}$	$4.3367896302 \times 10^{-17}$	$-4.18721614389 \times 10^{-16}$
22	$1.1603142 \times 10^{-21}$	$-1.24992369 \times 10^{-20}$	$1.356164500 \times 10^{-19}$	$-1.4385431446 \times 10^{-18}$

TABLE X: The case $\Lambda = -0.1$, $k = 0$: coefficients $A_{9,2i+1}$, $A_{10,2i+1}$, $A_{11,2i+1}$ and $A_{12,2i+1}$ of the polynomials v_9 , v_{10} , v_{11} and v_{12} . The coefficients of the normalization are $N_9 = 7.5849746655368418037$, $N_{10} = 8.0298788484656264819$, $N_{11} = 8.4547979707038394024$ and $N_{12} = 8.8624524914410403455$.

i	Coefficients $A_{13,2i+1}$	Coefficients $A_{14,2i+1}$	Coefficients $A_{15,2i+1}$	Coefficients $A_{16,2i+1}$
0	1	1	1	1
1	-28.874970949975714687	-32.223233211364481443	-35.665532275682033887	-39.198251529003849358
2	227.37032148479115396	286.18322037516669316	353.66043745913659654	430.30233892012025033
3	-763.33867329688880573	-1098.9702787419893209	-1533.3090062044208992	-2084.0733056378894925
4	1310.9566981411351041	2201.1873227411010990	3518.9507974024094019	5405.2859712063233703
5	-1253.2820823144540218	-2526.1788310714302717	-4723.4798592947973963	-8326.4852691380756713
6	682.8659047522031438	1733.7148182448332023	3910.3445264112686706	8082.2732257288263686
7	-201.42205297409225858	-710.5820774758294583	-2036.818888939089888	-5098.7078988356139768
8	23.83388279078582876	161.51383216003876259	656.7580090538630533	2104.4142883962174142
9	1.9964983432239179802	-13.983925825435929072	-120.01071532536019751	-554.29995868121941466
10	-0.6685900667196059490	-1.5243865249756624550	8.0886313897954242944	84.782128632037917752
11	-0.00526324112214527088	0.34761589239484017198	0.9591113608689540978	-4.8518135024721145179
12	0.00844311594833368901	0.00454698488229800367	-0.17490412409387389962	-0.52275410295941520127
13	0.000174146731635146709	-0.003485249000714049383	-0.001822245416712193038	0.086297493208029065181
14	-0.0000598451562116688980	-0.0000582113886434985875	0.001387134795031486997	0.000166148784298836630
15	-0.0000036802644318467220	0.0000198905161243789736	0.0000110751825193118621	-0.0005283014687449857168
16	0.00000008948000268032456	0.00000095390264285024449	-0.00000643970333012891868	0.00000216683354086455815
17	0.000000018958885951621149	-0.000000032423648042776966	-0.00000020634768304863584	0.00000196885695859247188
18	0.00000000979447180408524	-0.000000004317648229337035	0.000000011114789685221789	0.000000030318886497998249
19	$2.71616246788358 \times 10^{-11}$	-0.0000000001695948939628262	0.0000000008957869947573868	-0.00000000033547352564528893
20	$4.4018625809635 \times 10^{-13}$	$-3.42516125912610 \times 10^{-12}$	$2.471846771297929 \times 10^{-11}$	-0.00000000015941971769777072
21	$3.946485006354 \times 10^{-15}$	$-3.6168970617118 \times 10^{-14}$	$3.2067348095811 \times 10^{-13}$	$-2.72454248921859 \times 10^{-12}$
22	$1.5215597395 \times 10^{-17}$	$-1.58872844062 \times 10^{-16}$	$1.647260329029 \times 10^{-15}$	$-1.6927140309565 \times 10^{-14}$

TABLE XI: The case $\Lambda = -0.1$, $k = 0$: coefficients $A_{13,2i+1}$, $A_{14,2i+1}$, $A_{15,2i+1}$ and $A_{16,2i+1}$ of the polynomials v_{13} , v_{14} , v_{15} and v_{16} . The coefficients of the normalization are $N_{13} = 9.2549982289601205306$, $N_{14} = 9.6341574144155898259$, $N_{15} = 10.001394801227172017$ and $N_{16} = 10.357816917312201220$.

i	Coefficients $A_{17,2i+1}$	Coefficients $A_{18,2i+1}$	Coefficients $A_{19,2i+1}$	Coefficients $A_{20,2i+1}$
0	1	1	1	1
1	-42.818075477014044783	-46.522102414535798738	-50.307613353972484262	-54.172187839658743670
2	516.60124996673597985	613.04593626399226781	720.11780439718317891	838.29525508507567610
3	-2770.5184363380461860	-3613.4869945862085907	-4635.3958915081377870	-5860.2905303705903531
4	8031.0643052243792793	11600.594264758079964	16355.134985764342392	22576.823483403682361
5	-13992.376573406155088	-22597.400379554647521	-35285.868470667908997	-53526.528942521508014
6	15606.322037343566342	28522.411642485439145	49802.769083156847083	83665.654138659815313
7	-11565.353398244823011	-24306.842661588143797	-48040.616179513372979	-90235.832590244279443
8	5797.6407166951520743	14328.921222493045042	32574.308067403153125	69242.534529065824217
9	-1963.6315119914747301	-5902.0851094217960356	-15779.499229730758815	-38564.105889499156759
10	436.58402326206084665	1689.3844119449828328	5490.0412139413411117	15763.407929925976297
11	-58.096388332677072945	-326.37798591789254472	-1361.9469873825081100	-4741.6864982323540531
12	3.1114691105151574600	39.156744861307006175	234.54490257155531852	1041.9112893863402300
13	0.24603670167500815848	-2.1265929306784884977	-26.162891383355434394	-163.51438193058849124
14	-0.041701170848019458659	-0.092236589423893940527	1.5090010856945010152	17.379571503364081960
15	0.0004289240456083554886	0.019385890416302531676	0.016469307127428777114	-1.0812496755700901569
16	0.0001871920167131190035	-0.00047635782051305835602	-0.0083116101420201682610	0.015352271877295341103
17	-0.00000352657937019769039	-0.00005778228812711137480	0.00033924590352543960700	0.0029564870960007465462
18	-0.000000537367361145320176	0.000002184203662011836134	0.000012818089019123537845	-0.00018811565318134519679
19	0.000000008642339275576356	0.0000001165444563521529706	-0.000000949882865002101060	0.000000069245419364805160
20	0.000000008408248128099117	-0.0000000025438197808695685	-0.0000000130170676855048615	0.0000002867032868598201734
21	$2.181633244226363 \times 10^{-11}$	-0.00000000015901626316181483	0.00000000096318855304765349	-0.00000000314072046155475736
22	$1.72645172267161 \times 10^{-13}$	$-1.747765459211155 \times 10^{-12}$	$1.7570686102305507 \times 10^{-11}$	-0.000000000175470925339182865

TABLE XII: The case $\Lambda = -0.1$, $k = 0$: coefficients $A_{17,2i+1}$, $A_{18,2i+1}$, $A_{19,2i+1}$ and $A_{20,2i+1}$ of the polynomials v_{17} , v_{18} , v_{19} and v_{20} . The coefficients of the normalization are $N_{17} = 10.704554852432789228$, $N_{18} = 11.042302467661782828$, $N_{19} = 11.371947653621139153$ and $N_{20} = 11.694019311542212328$.

i	Coefficients $A_{21,2i+1}$	Coefficients $A_{22,2i+1}$	Coefficients $A_{23,2i+1}$
0	1	1	1
1	-58.113569811722124272	-62.129694188819134863	-66.218645891923138150
2	968.05133093681225903	1109.8555369654760655	1264.1735619567930545
3	-7313.8405815021465148	-9023.3742844288512616	-11017.892411056715348
4	30592.495945883842914	40777.824146671337701	53561.494870460181802
5	-79175.520459550748409	-114547.41165151219911	-162494.05150986880322
6	135962.25131813175913	214649.82636538022037	330362.32813557261476
7	-162347.07505899158433	-281466.38986469103875	-472495.16669313862312
8	139230.03723888623362	267101.52362763513657	492106.02976151703023
9	-87696.473111867246390	-187884.02174700301995	-382727.20362215702606
10	41166.134120203320700	99628.255907925668111	226391.61303713154891
11	-14516.830038316926307	-40270.402807365472386	-103199.15068593732476
12	3851.1192429595082125	12483.503210788413188	36573.960225972353470
13	-763.73249518978900371	-2970.9518664472614691	-10127.942085143099820
14	111.27157471232642190	540.33405521327578122	2194.2680274652986640
15	-11.473101658886635365	-74.197521867395963823	-370.97039963510504406
16	0.76784641910736935647	7.5117960938887405681	48.589506561001151270
17	-0.024795179287515729020	-0.53522777634520222233	-4.8664190536731115201
18	-0.00051868350247051926374	0.024126121954098631408	0.36475367507306886888
19	0.000078280717165341609642	-0.00045547223241067068401	-0.019759071102718174400
20	-0.0000019864464247742925383	-0.000014084210433377756998	0.00072912093021448666714
21	-0.00000003439787018249709564	0.00000099826862799863224994	-0.000016369627547200697053
22	0.000000001741290030126822791	-0.000000017175604877952762693	0.00000016843934455243322889

TABLE XIII: The case $\Lambda = -0.1$, $k = 0$: coefficients $A_{21,2i+1}$, $A_{22,2i+1}$ and $A_{23,2i+1}$ of the polynomials v_{21} , v_{22} and v_{23} . The coefficients of the normalization are $N_{21} = 12.009134872819683019$, $N_{22} = 12.317752618945874001$ and $N_{23} = 12.619905741748600855$.

i	Coefficients $A_{1,2i+1}$	Coefficients $A_{2,2i+1}$	Coefficients $A_{3,2i+1}$	Coefficients $A_{4,2i+1}$
0	1	1	1	1
1	21.8377842950992	18.4334293031812	15.1810746656932	12.0920975305193
2	147.651955803377	105.526579144947	71.7770440011361	45.5993383732682
3	483.861595247111	292.175608570147	163.214861374083	81.5591541074805
4	931.239526525682	471.973074791482	213.571689647027	81.2934444080791
5	1170.31529375274	492.198865923421	176.051457215035	47.8082621918444
6	1026.20921841086	351.941764334297	95.3901450241527	15.8144829846098
7	656.457818963428	178.839126369668	33.9450844505980	1.75908152275731
8	316.065578003045	65.6921984104879	7.27484182555577	-0.791910624911047
9	117.125256376888	17.3696744876673	0.491066910363862	-0.402967416824580
10	33.9440610584924	3.16006349470625	-0.229720629183342	-0.0768400028663149
11	7.77907712686178	0.324132553137970	-0.0886170587898875	-0.00292105263872861
12	1.41970868793207	-0.00915418415050491	-0.0154236544033143	0.00185443062508932
13	0.207044911107463	-0.0106898237097331	-0.00126626701898463	0.000452251846104077
14	0.0241257915458056	-0.00227263356521181	0.000050915914977872	0.0000453164274436939
15	0.00223753092789531	-0.000300778426490677	0.0000295746638122252	0.00000034783472614020
16	0.000163799213077448	-0.0000282611500985331	0.00000437665019606404	-0.000000510363960204255
17	0.0000093321070409920	-0.00000194789534022172	0.000000393985207700712	-0.0000000746297051171108
18	0.00000040460640223067	-0.0000000983927199169808	0.0000000240272842869963	-0.00000000586769020323061
19	0.0000000128813277100826	-0.00000000355572877567576	0.00000000100402369449065	-0.000000000291275827943990
20	0.0000000002836090970636	-8.7189990778683 $\times 10^{-11}$	2.76996707360698 $\times 10^{-11}$	-9.1566862013583 $\times 10^{-12}$
21	3.854306040188 $\times 10^{-12}$	-1.3007318907487 $\times 10^{-12}$	4.5621367925570 $\times 10^{-13}$	-1.67542869114651 $\times 10^{-13}$
22	2.43401031613 $\times 10^{-14}$	-8.91495847348 $\times 10^{-15}$	3.4044147473993 $\times 10^{-15}$	-1.3653666863045 $\times 10^{-15}$

TABLE XIV: The case $\Lambda = -0.1$, $k = -1$: coefficients $A_{1,2i+1}$, $A_{2,2i+1}$, $A_{3,2i+1}$ and $A_{4,2i+1}$ of the polinomys v_1 , v_2 , v_3 and v_4 . The coefficients of the normalization are $N_1 = 0.107994257450045 \times 10^{-6}$, $N_2 = 0.292690588075209 \times 10^{-5}$, $N_3 = 0.518252990768902 \times 10^{-4}$ and $N_4 = 0.683649822517353 \times 10^{-3}$.

i	Coefficients $A_{5,2i+1}$	Coefficients $A_{6,2i+1}$	Coefficients $A_{7,2i+1}$	Coefficients $A_{8,2i+1}$
0	1	1	1	1
1	9.18190533268822	6.47301073960000	4.00115813990478	1.80436900132522
2	26.1743762082396	12.6594902235932	4.16903822033216	-0.299804601981924
3	34.0957320592592	9.92492684765362	0.26293559391319	-1.64081279306699
4	22.4752715608062	2.00505672302545	-1.74796349658398	-0.510193340358966
5	6.37807566817418	-1.67761640342591	-0.827957567330092	0.306140893014826
6	-0.805445298790648	-1.1118485823729	0.0481471210374753	0.175238091233888
7	-1.19296920822983	-0.168747412574362	0.123505645725467	-0.0023207172666482
8	-0.343569927303758	0.0577809210608984	0.0248551679945839	-0.0173518187572286
9	-0.0165422158611707	0.0266126276608532	-0.00373900679066606	-0.00275760349002051
10	0.0145426929549039	0.00263861673458369	-0.00204171593286728	0.000531936179331518
11	0.00396245317708414	-0.000613920870654406	-0.000175415857599938	0.000189123010906910
12	0.000311820881878200	-0.000192175631525371	0.0000411479287477862	0.00000625079217426494
13	-0.0000461908583584600	-0.0000140755743974023	0.0000100324849830199	-0.00000413971266272626
14	-0.0000137611226075986	0.00000189045389193022	0.000000405268756113207	-0.000000569328117910824
15	-0.00000129418233580770	0.000000468942461721557	-0.000000113104785867229	0.00000000893632522180967
16	-0.0000000040791859495057	0.0000000327343587403344	-0.0000000170097722403145	0.00000000790985805873956
17	0.0000000119786522592102	-0.000000000881155828875187	-0.000000000590109758554436	0.000000000608101054536502
18	0.00000000142024608946527	-0.00000000033392082180239	7.09820188285488 $\times 10^{-11}$	-8.08849678866289 $\times 10^{-12}$
19	8.74468304111290 $\times 10^{-11}$	-2.75569519071327 $\times 10^{-11}$	9.46108114535898 $\times 10^{-12}$	-3.99957955781206 $\times 10^{-12}$
20	3.18375051718700 $\times 10^{-12}$	-1.18697762945545 $\times 10^{-12}$	4.95934160671152 $\times 10^{-13}$	-2.65204337531615 $\times 10^{-13}$
21	6.51558566491249 $\times 10^{-14}$	-2.73576846957429 $\times 10^{-14}$	1.29509232785582 $\times 10^{-14}$	-7.87801182225090 $\times 10^{-15}$
22	5.81401010856087 $\times 10^{-16}$	-2.67800465237542 $\times 10^{-16}$	1.39083409062693 $\times 10^{-16}$	-9.25938745797130 $\times 10^{-17}$

TABLE XV: The case $\Lambda = -0.1$, $k = -1$: coefficients $A_{5,2i+1}$, $A_{6,2i+1}$, $A_{7,2i+1}$ and $A_{8,2i+1}$ of the polinomys v_5 , v_6 , v_7 and v_8 . The coefficients of the normalization are $N_5 = 0.699677173459493 \times 10^{-2}$, $N_6 = 0.556647587279093 \times 10^{-1}$, $N_7 = 0.330418318317813$ and $N_8 = 1.29283805747619$.

l	Coefficients $A_{9,2i+1}$	Coefficients $A_{10,2i+1}$	Coefficients $A_{11,2i+1}$	Coefficients $A_{12,2i+1}$
0	1	1	1	1
1	-0.232010916759383	-2.43487417097320	-4.93455597590118	-7.69645852094990
2	-1.85623104021040	-0.73836052792331	4.05659416695777	13.7141424722470
3	-0.265450448980053	2.04821420478302	2.31824860373187	-4.94408507267326
4	0.786253245370039	0.420288826610130	-2.23653053543589	-4.28380968001096
5	0.232581678867701	-0.546742403426884	-0.530835364516564	2.28767539275548
6	-0.107274956602857	-0.130901720787139	0.424516857735247	0.562747928024902
7	-0.0481118639375175	0.0598259416409346	0.081622419872838	-0.333454264677029
8	0.00335207257698909	0.0184283915885142	-0.0375565130712911	-0.0479110227923248
9	0.00390684508813047	-0.00256898389904225	-0.00776250127887547	0.0239025167554597
10	0.000293677739112035	-0.00124048844412143	0.00162742947212471	0.00298919015302213
11	-0.000131500602009393	0.0000006208725404758	0.000420150252596240	-0.000934483564273605
12	-0.0000234385715011634	0.0000407506272747854	-0.000029049828816688	-0.000126408733730916
13	0.00000108202211165345	0.00000321248157945464	-0.0000124807815871202	0.0000196909777083217
14	0.000000573039490535006	-0.000000580085543479874	-0.000000127374065571837	0.00000327880652990617
15	0.0000000324470400434446	-0.0000000929895742739634	0.000000190970152308110	-0.000000186559743938849
16	-0.00000000446711096592629	0.000000000765718977012007	0.0000000121032382246325	-0.0000000480975160590495
17	-0.000000000659364775626084	0.000000000948169589019447	-0.00000000119161228215810	-0.00000000011776394937123
18	$-1.55485486252239 \times 10^{-11}$	$5.47596862462946 \times 10^{-11}$	$-0.000000000151948575638067$	$0.000000000350699137653205$
19	$2.55875406226149 \times 10^{-12}$	$-1.89533455884707 \times 10^{-12}$	$-1.16945920999965 \times 10^{-12}$	$1.46509482512341 \times 10^{-11}$
20	$2.33518330702113 \times 10^{-13}$	$-3.30715164626864 \times 10^{-13}$	$5.49785532554198 \times 10^{-13}$	$-8.22408951431525 \times 10^{-13}$
21	$7.98100028352004 \times 10^{-15}$	$-1.35925861467088 \times 10^{-14}$	$2.98035371962606 \times 10^{-14}$	$-7.21928400402298 \times 10^{-14}$
22	$1.02803953739253 \times 10^{-16}$	$-1.95362588495032 \times 10^{-16}$	$4.93162483077208 \times 10^{-16}$	$-1.43434382739745 \times 10^{-15}$

TABLE XVI: The case $\Lambda = -0.1$, $k = -1$: coefficients $A_{9,2i+1}$, $A_{10,2i+1}$, $A_{11,2i+1}$ and $A_{12,2i+1}$ of the polynomials v_9 , v_{10} , v_{11} and v_{12} . The coefficients of the normalization are $N_9 = 2.92703125957288$, $N_{10} = 4.35635774997061$, $N_{11} = 5.39681832616858$ and $N_{12} = 6.24220211798654$.

i	Coefficients $A_{13,2i+1}$	Coefficients $A_{14,2i+1}$	Coefficients $A_{15,2i+1}$	Coefficients $A_{16,2i+1}$
0	1	1	1	1
1	-10.6841664013405	-13.8742150817412	-17.2503270722203	-20.7999881540638
2	29.3147115073451	51.8840233684938	82.4201447999443	121.901220187412
3	-27.8107960004119	-76.9972681817258	-165.993521308318	-311.189756974583
4	4.6218685711793	46.6058507383930	160.587831324991	407.968109284316
5	6.17023638414212	-3.5351498009348	-69.8936934649346	-287.865946240682
6	-2.25304381758315	-7.59830396617375	2.86651578722247	99.0775488526631
7	-0.50834042972362	2.21102625218132	8.34823080918794	-4.08633289155962
8	0.260659211268764	0.389521458880789	-2.19601372488250	-8.30119517836824
9	0.0233981838235592	-0.200572813222270	-0.236668045675840	2.18686271337729
10	-0.0148903861397447	-0.0066975973688412	0.148725930312016	0.0783477485345212
11	-0.000821801814455150	0.00878058536092591	-0.00324654886311030	-0.102429805871289
12	0.000488171433380495	-0.000056446659015478	-0.00466781743553869	0.00774541374029079
13	0.0000253800980710251	-0.000224658192818058	0.000304723373636126	0.00203081309779073
14	-0.00000948811922148724	0.0000031236965216455	0.0000840098760734869	-0.000278333835922816
15	-0.000000584760574386555	0.00000356209497246444	-0.00000678634432331948	-0.0000194091549503491
16	0.000000104804340274713	-0.0000000235206434466662	-0.000000943013330726463	0.00000413943671070284
17	0.00000000824331415372495	-0.0000000345784393018707	0.0000000733068104143618	0.000000078877544486722
18	-0.000000000555486232727475	-0.00000000024777655854987	0.00000000674723462090878	-0.0000000320870558331197
19	$-6.10779307199931 \times 10^{-11}$	$0.000000000187683405960227$	$-0.000000000397367068213861$	$-1.5160047897411 \times 10^{-11}$
20	$4.69013712315356 \times 10^{-13}$	$4.18930259721944 \times 10^{-12}$	$-2.88588907307502 \times 10^{-11}$	$0.000000000128509091870770$
21	$1.80184547774889 \times 10^{-13}$	$-4.33041646241597 \times 10^{-13}$	$8.66474144450464 \times 10^{-13}$	$-5.1077434842036 \times 10^{-13}$
22	$4.5727033338865 \times 10^{-15}$	$-1.56130759518453 \times 10^{-14}$	$5.63055942629174 \times 10^{-14}$	$-2.12450030498590 \times 10^{-13}$

TABLE XVII: The case $\Lambda = -0.1$, $k = -1$: coefficients $A_{13,2i+1}$, $A_{14,2i+1}$, $A_{15,2i+1}$ and $A_{16,2i+1}$ of the polynomials v_{13} , v_{14} , v_{15} and v_{16} . The coefficients of the normalization are $N_{13} = 6.98413261507519$, $N_{14} = 7.65951898965145$, $N_{15} = 8.28713472086337$ and $N_{16} = 8.87811017222452$.

i	Coefficients $A_{17,2i+1}$	Coefficients $A_{18,2i+1}$
0	1	1
1	-24.5135934862227	-28.3826830252342
2	171.297635578830	231.563657333425
3	-532.032302271353	-851.104089577650
4	879.049849725663	1700.00285906748
5	-837.566899410330	-2028.92845736494
6	466.195823019623	1512.52404634436
7	-137.342049715716	-704.222275510342
8	8.59568426834457	188.892562917903
9	7.36166712174031	-17.5315261277321
10	-2.12487622903933	-5.41896929313905
11	0.0616353109016188	1.93593317266994
12	0.0612186597075610	-0.163977083115011
13	-0.00828671964266124	-0.0263879773884429
14	-0.00051214720454678	0.00637464986776314
15	0.000167433511984800	-0.000186792886772932
16	-0.00000303760478624729	-0.0000646259429141273
17	-0.00000147924868677497	0.00000596587830124041
18	0.000000075304283408867	0.000000160392422977485
19	0.00000000606586647233395	-0.0000000400489171784274
20	-0.000000000434054633266702	0.00000000080165107135159
21	$-9.3095562785296 \times 10^{-12}$	$8.54942653394270 \times 10^{-11}$
22	$8.32964788661547 \times 10^{-13}$	$-3.37515341995877 \times 10^{-12}$

TABLE XVIII: The case $\Lambda = -0.1$, $k = -1$: coefficients $A_{17,2i+1}$ and $A_{18,2i+1}$ of the polinomys v_{17} and v_{18} . The coefficients of the normalization are $N_{17} = 9.43144441465169$ and $N_{18} = 9.75064661119335$.




FIG. 1: Behavior of the expectation value of the scalar factor for $\Lambda = -0.1$, $k = 1$.




FIG. 2: Behavior of the expectation value of the scalar factor for $\Lambda = -0.1$, $k = 0$.




FIG. 3: Behavior of the expectation value of the scalar factor for $\Lambda = -0.1$, $k = -1$.